How The Pest Was Won

Dr G Percival: Bartlett Tree Research Laboratory



Bartlett Tree Research Laboratory, University of Reading

Pest and Diseases Within the UK: Increased severity





Pests and Diseases over the past 10 years



Ever Increasing Threats



Chalara ash die-back estimated to kill 99% of all UK ash trees



Oak Processionary Moth-Dangerous to humans

FAGACEAE 000-73-193

1413


"TURKEY OAK" QUERCUS CERRIS

S.EUROPE & W.ASIA

General Prediction

Problem will get worse

What are the Options?





P.ramorum 3,000,000 larch trees felled....no effect on spread



Thankfully there is a different approach



SPECIATOR 10 NOVEMBER 2012 I WWW SPECTATOR

What are the Options







A Different Approach

Western medicine dictates that preventation of infectious diseases (typhoid, diphtheria, measles, hepatitis, small pox) is primarily via vaccination. In such circumstances the human body is injected with a weakened strain of a disease. This in turn stimulates the body to produce antibodies against that specific disease which in turn confers immunity. Importantly a "one-off" vaccination can confer immunity for many years (at least 10) and in some cases last an entire life time.





Can we use these vaccination principles for trees?

The answer is yes. Vaccinating plants against pests and diseases is not a new concept; the idea of inducing resistance in response to plant diseases was recognised in the early 20th century when heat or cold treated *Botrytis cinerea* (grey mould) when exposed to *Begonia* plants instead of causing infection as expected, resulted in the plants developing resistance.



Several studies have found that "vaccinating" trees to be effective in controlling:

- Fire blight (Erwinia carotovora)
- Phytophthora root rot.



- Powdery mildew (Sphaerotheca pannosa var. rosa, Phyllactinia sp and Uncinula necator)
- Wilt disease of spruce (Ceratocystis polonica)
- Importantly, the level of disease control achieved was comparable with currently used agrochemicals and a "oneoff" vaccination has been shown to provide growing season protection.

- Interestingly tree defence responses are superior to that of humans!
- An injection against typhoid would only confer immunity against typhoid. Further separate injections would be required if immunity against diphtheria or measles was required.
- In trees, however, a single vaccination causes:
- Accumulation of antimicrobial proteins, fungi-toxic enzymes, phenolics and terpenoids within leaves, stems and roots.
- Leaves become thicker and more lignified.
- Enhanced resin production, production of phenolics and initiation of a wound periderm occurs.
- Importantly, because multiple defence mechanisms are switched on it is it highly unlikely that pests and diseases can develop resistance to this measure.
- In addition, a single vaccination has been shown to provide resistance against biologically different pathogens (fungal, bacterial, virus) over a growing season

A small but significant step.

- Trees can be vaccinated by applying products as a root drench! (Percival G.C and Banks J M (2015). Arboricultural Journal: 37(1): 7-20
- Applying products via the roots opens up opportunities to manage tree pest and diseases without the need to sbpray.









So what soil amendments can we use to vaccinate trees?

▶ 1. A pure mulch.





Willow mulch





Willow mulch



Provides a suitable soil environment for root growth and mycorrhizal associations.



Slide courtesy Drs D Zwart/K Fite

Phytophthora Management

Vinca and Gardenia inoculated with Phytophthora Control Compost Biochar



Biochar effect on leaf blotch and leaf miner severity



Biochar 0.50kg m²

Biochar 0.25kg m²

Phosphites







Chitin – What is it?

2nd most widespread natural polymer

- Forms structure of:
 - Fungi cell walls
 - Insect exoskeletons
 - Crustacean exoskeletons
- Insoluble!
 - Derivatives soluble... and more effective

S S

Apple scab trial site

Control





R&D Trials Over the Past 2 Years

Looking at these four products singly and in combination i.e. Biochar Biochar + Chitin Biochar + Pure Mulch Biochar + Phosphites Biochar + Chitin + Pure Mulch etc.

Importantly Biochar, Mulch, Chitin are waste products.



Woodland/Forest



Business Park





2000 trees in an urban landscape setting




Barnet Council Trial Month 4 After Planting Street Plantings







Back Site has a unique set of pests and diseases



Ash Die-back



Scab and Blister Mites

Is it Working? Rust Disease







HCLM and Pseudomonas Bleeding Canker





Pi + Chit + Biochar + Mulch

Control






Defoliating beetles




Honey Fungus (*Armillaria* spp)




Specific activity of defensive root enzymatic activity Vs ARMILLARIA severity

Disease severity




Specific activity (SOD)

Specific activity of defensive root enzymatic activity Vs ARMILLARIA severity




Thicker leaves



treatment 0.25% fracture 050.tif

Control

Biochar + Chitin



Summary

Use of biochar, pure mulch, chitin and phosphites do:

1. Cause enhancement of defensive enzymes in leaves and roots (SOD, Peroxidase, Beta Glucanase)
2. Leaves become thicker and more lignified.
3. Highly likely many more defense enzymes/metabolites enhanced that we haven't analysed for.

►4. 15-20% reduction in growth

But it's not perfect

- Using this system can provide reductions of disease/pest severity ranging from 30-70%.
- Using conventional plant protection products can provide 100% reductions.



► For the future: Willow species



► For the future; Chitin and/or Chitosan



► For the future; Biochar



Practical Guidelines:

- Mulch: Willow; 5-10 cm deep, mulch area under crown + 1 metre beyond
- Chitin: 120g per square metre
- Phosphites: 1.5 litres per 100 litres of water
- Biochar: 5% by soil volume i.e. for every 19 litres of soil add 1 litre of biochar.
- How long does the
 induced response last'
- ▶ ?????





END